



Cosmetic Vein Centers of Virginia

RECOMMENDATIONS FOR BETTER LEGS

REGULAR EXERCISE

Walking, running, stairmaster, aerobics, swimming, elliptical machine, or biking for 30 minutes, 5-7 days per week will help reduce aching, pain and tiredness in your legs.

ELEVATE YOUR LEGS

Elevating your legs above heart level for at least 10 minutes once or twice daily may diminish aching and swelling.

MOVE YOUR LEGS FREQUENTLY

Flexing your ankles 10 times will pump the blood out of your legs like walking does. Repeat this every 10 minutes while standing or sitting and try to walk for at least 2 minutes every half hour.

AVOID WEARING HIGH HEELS

Wearing high heels interferes with the normal pumping action that occurs when you walk and may lead to aching and cramping of the legs.

MAINTAIN A PROPER WEIGHT

Even moderate weight loss may reduce aching in the legs due to varicose veins and diminish the rate at which spider veins develop.

WEAR SUPPORT HOSE

Available at pharmacies and medical supply stores. There are many brands to choose from. Lighter support stockings are available at department stores. However, it is best to wear a stocking that is labeled "graduated" as this will really help to improve your vein function.

Light support: 4-14 mm

Moderate support : 15-20 mm

Heavy support : 20-30 mm

Prescription strength: 30-40 mm or above

David T. MacMillan, M.D., FACS

1901 South Main St., Suite 2
Blacksburg, VA 24060
Phone 540-552-VEIN (8346)
Fax 540-951-VEIN (8346)
www.veinrx.com